

On The Henstock- Kurzweil Integral For Riesz-Spaces-Valued Functions Defined On Euclidean Space \mathfrak{R}^n

Muslim Ansori
Mathematics Departement, Universitas Lampung
Email : ansomath@yahoo.com

Abstract

This paper is a partial result of our researchs in the main topic "On The Henstock-Kurzweil Integral for Riesz-Spaces-valued Functions Defined on Riesz Space L ". We construct Henstock-Kurzweil integral for Riesz-spaces-valued functions defined on Euclidean space \mathfrak{R}^n and prove some basic properties among which the fact that our new integral is coincides with the Henstock-Kurzweil Integral for Banach-spaces valued functions defined on space \mathfrak{R}^n .

Keywords : Riesz Space, Henstock-Kurzweil Integral

1. INTRODUCTION

The Henstock-Kurzweil integral for Riesz-space-valued functions defined on bounded subintervals of the real line and with respect to operator-valued measures was investigated by Riecan(1989,1992) and Riecan and Brabelova(1996), with respect to (D) - convergence (that is a kind of convergence in which the ε -technique is replaced by a technique involving double sequences, see Riecan and Neubrunn(1997)), with respect to the order convergence, see Boccuto(1998) and in Boccuto and Riecan(2004) with respect to the order convergence but the Henstock-Kurzweil integral for Riesz-space-valued functions was defined on unbounded subintervals of the real line.

The Henstock-Kurzweil integral for real-valued functions defined on Euclidean space \mathfrak{R}^n with respect to volume α was investigated in Pfeffer(1993) and Indrati(2002) and The Henstock-Kurzweil integral for bounded-sequence-space-valued functions defined on Euclidean space \mathfrak{R}^n with respect to volume α was investigated in Muslim and Soeparna(2002) and Zachriwan(2004).

The main goal of this paper is to generalize the results above by constructing Henstock-Kurzweil integral for Riesz-valued functions defined on Euclidean space \mathfrak{R}^n and we prove some fundamental properties.

2. PRELIMINARY

Let \mathbb{N} be the set of all strictly positive integers, \mathfrak{R} the set of the real numbers, \mathfrak{R}^+ be the set of all strictly positive real numbers. Moreover, we refer to (Pfeffer,1993)

about the notions of cell, segmentation, partition, α -volume, and δ -fine Perron partition.

Definisi 2.1 (Zaanen,1996) : A Riesz space L is said to be Dedekind complete if every nonempty subset of L , bounded from above, has supremum in L .

Definisi 2.2 (Riecan, 1998) : A bonded double sequence $(a_{i,j})_{i,j} \in L$ is called regulator or (D) -sequence if, for each $i \in \mathbb{N}$, $a_{i,j} \downarrow 0$, that is $a_{i,j} \geq a_{i,j+1} \forall j \in \mathbb{N}$ and $\bigwedge_{j \in \mathbb{N}} a_{i,j} = 0$.

Definisi 2.3 (Boccuto and Riecan, 2004) : Given a sequence $(r_n)_n \in L$. Sequence $(r_n)_n$ is said to be (D) -convergence to an element $r \in L$ if there exist a regulator $(a_{i,j})_{i,j}$, satisfying the following condition:

for every mapping $\rho : \mathbb{N} \rightarrow \mathbb{N}$, there exists an integer n_0 sehingga $|r_n - r| \leq \bigvee_{i=1}^{\infty} a_{i,\rho(i)}$ for all $n \geq n_0$. In this case, the notation is denoted by $(D)\lim_n r_n = r$.

Definition 2.4 (Boccuto and Riecan, 2004) : A Riesz Space L is said to be weakly σ -distributive if for every (D) - sequence $(a_{i,j})$, then

$$\bigwedge_{\rho \in \mathbb{N}^{\mathbb{N}}} \left(\bigvee_{i=1}^{\infty} a_{i,\rho(i)} \right) = 0.$$

Throughout the paper, we shall always assume that L is Dedekind complete weakly σ -distributive Riesz space.

Main Results

In the principle, this integral is a generalization of Henstock-Kurzweil integral for Riesz-valued functions defined on subintervals of the real line by changing the length of $[a,b] \subset \mathbb{R}$ with the general volume α of a cell $A \subset \mathbb{R}^n$. See **Pfeffer**(1993) and **Muslim and Soeparna**(2002). Remember that the volume α on cell $A \subset \mathbb{R}^n$ is an additive and non negative function from $\mathfrak{T}(A)$ into \mathbb{R} , where $\mathfrak{T}(A)$ is a collection of all subcells in A .

Definition 3.1 : Let α be a volume on \mathfrak{R}^n and $A \subset \mathfrak{R}^n$ be a cell. A function $f : \mathfrak{R}^n \rightarrow L$ is said to be Henstock-Kurzweil integrable on A with respect to α , denoted by $f \in HK(A, L, \alpha)$, if there exists an element $\Xi \in L$ and (D) -sequence $(a_{i,j})_{i,j} \in L$ such that for every $\rho \in \square^\square$ we can find a function $\delta : E \rightarrow \mathfrak{R}^+$ such that

$$\left| P \sum f(\bar{x}) \alpha(I) - \Xi \right| = \left| \sum_{k=1}^r f(\bar{x}_k) \alpha(I_r) - \Xi \right| \leq \bigvee_{i=1}^{\infty} a_{i, \rho(i)}$$

for every δ -fine Perron partition $P = \{(I, \bar{x})\} = \{(I_1, \bar{x}_1), (I_2, \bar{x}_2), \dots, (I_r, \bar{x}_r)\}$ on A .

We note that the Henstock-Kurzweil integral with respect to α is well- defined, that is there exists at most one element Ξ , satisfying Definition 3.1 and in this case we have $(HK) \int_A f d\alpha = \Xi$. The uniqueness is given in the following theorem.

Theorem 3.2 : Let α be a volume on \mathfrak{R}^n and $A \subset \mathfrak{R}^n$ be a cell. If function $f \in HK(A, L, \alpha)$, then its α -integral is unique.

Proof: Let $f \in HK(A, L, \alpha)$. If both Ξ_1 and Ξ_2 are Henstock-Kurzweil integral of function f , satisfying Definition 3.1, then there exists two (D) -sequence $(a_{i,j})_{i,j}$ and $(b_{i,j})_{i,j}$ in L such that for every $\rho \in \square^\square$, we can find two positive function δ_1 and δ_2 on A , respectively, and for every δ_1 -fine Perron partition $P_1 = \{(I, \bar{x})\}$ and δ_2 -fine Perron partition $P_2 = \{(I, \bar{x})\}$ on A , we have

$$\left| P_1 \sum f(\bar{x}) \alpha(I) - \Xi_1 \right| \leq \bigvee_{i=1}^{\infty} a_{i, \rho(i)}$$

and

$$\left| P_2 \sum f(\bar{x}) \alpha(I) - \Xi_2 \right| \leq \bigvee_{i=1}^{\infty} b_{i, \rho(i)}$$

respectively. Let now $\delta(\bar{x}) = \min\{\delta_1(\bar{x}), \delta_2(\bar{x})\}$, for every $\bar{x} \in A$ and take any δ -fine Perron partition $P = \{(I, \bar{x})\}$ on A , then $P = \{(I, \bar{x})\}$ is both δ_1 -fine Perron partition and δ_2 -fine Perron partition on A , and thus we have

$$\begin{aligned}
 0 \leq |\Xi_1 - \Xi_2| &\leq \left| P_1 \sum f(\bar{x})\alpha(I) - \Xi_1 \right| + \left| P_2 \sum f(\bar{x})\alpha(I) - \Xi_2 \right| \leq \bigvee_{i=1}^{\infty} a_{i,\rho(i)} + \bigvee_{i=1}^{\infty} b_{i,\rho(i)} \\
 &\leq \bigvee_{i=1}^{\infty} (a_{i,\rho(i)} + b_{i,\rho(i)}) \\
 &\leq \bigvee_{i=1}^{\infty} c_{i,\rho(i)}
 \end{aligned}$$

where $c_{i,j} = 2(a_{i,j} + b_{i,j}) \quad \forall i, j \in \mathbb{N}$. By arbitrariness of $\rho \in \mathbb{N}^{\mathbb{N}}$, we get

$$0 \leq |\Xi_1 - \Xi_2| \leq \bigwedge_{\rho \in \mathbb{N}^{\mathbb{N}}} \left(\bigvee_{i=1}^{\infty} c_{i,\rho(i)} \right) = 0$$

since $c_{i,j}$ is (D) -sequence and thanks to weak σ -distributivity of L . Thus $\Xi_1 = \Xi_2$, and so our HK-integral is well-defined. \square

Now, we give some fundamental properties of $HK(A, L, \alpha)$.

Theorem 3.3 : If $f_1, f_2 \in HK(A, L, \alpha)$ and $k_1, k_2 \in \mathfrak{R}$, then $k_1 f_1 + k_2 f_2 \in HK(A, L, \alpha)$ and

$$(HK) \int_A (k_1 f_1 + k_2 f_2) d\alpha = k_1 (HK) \int_A f_1 d\alpha + k_2 (HK) \int_A f_2 d\alpha.$$

Proof : The proof is similar to the one of (Muslim, 2003), Theorem 3.1.3

Theorem 3.4 : If $f, g \in HK(A, L, \alpha)$ and $f(\bar{x}) \leq g(\bar{x})$ for every $\bar{x} \in A$, then

$$(HK) \int_A f d\alpha \leq (HK) \int_A g d\alpha.$$

Proof : By hypothesis, there exists two (D) -sequences, $(a_{i,j})_{i,j}$ and $(b_{i,j})_{i,j}$ such that, for every $\rho \in \mathbb{N}^{\mathbb{N}}$, we can find positive functions δ_1 dan δ_2 , respectively on A , and whenever $P_1 = \{(I, \bar{x})\}$ is δ_1 -fine Perron partition and $P_2 = \{(I, \bar{x})\}$ is δ_2 -fine Perron partition on A , we have

$$\begin{aligned}
 \left| P_1 \sum f(\bar{x})\alpha(I) - \int_A f d\alpha \right| &\leq \bigvee_{i=1}^{\infty} a_{i,\rho(i)} \Leftrightarrow \\
 \int_A f d\alpha - \bigvee_{i=1}^{\infty} a_{i,\rho(i)} &\leq P_1 \sum f(\bar{x})\alpha(I) \leq \int_A f d\alpha + \bigvee_{i=1}^{\infty} a_{i,\rho(i)}
 \end{aligned}$$

and

$$\left| P_2 \sum f(\bar{x}) \alpha(I) - \int_A g d\alpha \right| \leq \bigvee_{i=1}^{\infty} b_{i,\rho(i)} \Leftrightarrow$$

$$\int_A g d\alpha - \bigvee_{i=1}^{\infty} b_{i,\rho(i)} \leq P_2 \sum g(\bar{x}) \alpha(I) \leq \int_A g d\alpha + \bigvee_{i=1}^{\infty} b_{i,\rho(i)}$$

respectively.

For every $\bar{x} \in A$, let $\delta(\bar{x}) = \min\{\delta_1(\bar{x}), \delta_2(\bar{x})\}$, and take δ -fine Perron partition $P = \{(I, \bar{x})\}$ on A , then $P = \{(I, \bar{x})\}$ is both δ_i -fine Perron partition ($i = 1, 2$) on A . Thus we get

$$\int_A f d\alpha - \bigvee_{i=1}^{\infty} a_{i,\rho(i)} \leq P \sum f(\bar{x}) \alpha(I) \leq P \sum g(\bar{x}) \alpha(I) \leq \int_A g d\alpha + \bigvee_{i=1}^{\infty} b_{i,\rho(i)}$$

and hence, for every $\rho \in \square^\square$,

$$\int_A f d\alpha - \int_A g d\alpha \leq \bigvee_{i=1}^{\infty} a_{i,\rho(i)} + \bigvee_{i=1}^{\infty} b_{i,\rho(i)} \leq \bigvee_{i=1}^{\infty} c_{i,\rho(i)}$$

where $c_{i,j} = 2(a_{i,j} + b_{i,j}) \forall i, j \in \square$. By arbitrariness of $\rho \in \square^\square$, since $c_{i,j}$ is a (D) -sequence and taking into account of weak σ -distributivity of L , we get

$$\int_A f d\alpha - \int_A g d\alpha \leq \bigwedge_{\rho \in \square^\square} \left(\bigvee_{i=1}^{\infty} c_{i,\rho(i)} \right) = 0$$

that is $\int_A f d\alpha \leq \int_A g d\alpha$. This concludes the proof. \square

Definition 3.5 (Elementary Set): A set $A \subset \mathfrak{R}^n$ which is union of finite cells is called an elementary set.

Every elementary set can be segmented into non-overlapping cells. If A_1 and A_2 are elementary sets then $A_1 \cup A_2$ and $\overline{A_1 \setminus A_2}$ are also elementary sets. Integration on elementary set can be constructed through the following theorem.

Teorema 3.6 : Let α be a volume on \mathfrak{R}^n and A_1 and A_2 be non-overlapping cells in \mathfrak{R}^n and $A = A_1 \cup A_2$. If $f \in HK(A_1, L, \alpha)$ and $f \in HK(A_2, L, \alpha)$, then $f \in HK(A, L, \alpha)$ and

$$(HK) \int_{A=A_1 \cup A_2} f d\alpha = (HK) \int_{A_1} f d\alpha + (HK) \int_{A_2} f d\alpha$$

Proof : Let $f \in HK(A_1, L, \alpha)$ and $f \in HK(A_2, L, \alpha)$. There exists two (D) -sequence $(a_{i,j})_{i,j}$ and $(b_{i,j})_{i,j}$, such that for every $\rho \in \square^\square$, we can find positive functions δ_1 and

δ_2 on A respectively. Whenever $P_1 = \{(I, \bar{x})\}$ is δ_1 -fine Perron partition on A_1 and $P_2 = \{(I, \bar{x})\}$ is δ_2 -fine Perron partition on A_2 , we have

$$\left| P_1 \sum f(\bar{x}) \alpha(I) - \int_{A_1} f d\alpha \right| \leq \bigvee_{i=1}^{\infty} a_{i, \rho(i)}$$

and

$$\left| P_2 \sum f(\bar{x}) \alpha(I) - \int_{A_2} f d\alpha \right| \leq \bigvee_{i=1}^{\infty} b_{i, \rho(i)}$$

Let now $\delta : A \rightarrow \mathfrak{R}^+$ be such that,

$$\delta(\bar{x}) = \begin{cases} \delta_1(\bar{x}) & \text{if } \bar{x} \in A_1 \text{ and } \bar{x} \notin A_2 \\ \delta_2(\bar{x}) & \text{if } \bar{x} \in A_2 \text{ and } \bar{x} \notin A_1 \\ \min\{\delta_1(\bar{x}), \delta_2(\bar{x})\} & \text{if } \bar{x} \in A_1 \cap A_2 \end{cases}$$

for every δ -fine Perron partition $P = \{(I, \bar{x})\}$ on A where $P = P_1 \cup P_2$. Therefore, we get

$$\begin{aligned} & \left| P \sum f(\bar{x}) \alpha(I) - \left(\int_{A_1} f d\alpha + \int_{A_2} f d\alpha \right) \right| \\ & \leq \left| P_1 \sum f(\bar{x}) \alpha(I) - \int_{A_1} f d\alpha \right| + \left| P_2 \sum f(\bar{x}) \alpha(I) - \int_{A_2} f d\alpha \right| \\ & \leq \bigvee_{i=1}^{\infty} a_{i, \rho(i)} + \bigvee_{i=1}^{\infty} b_{i, \rho(i)} \leq \bigvee_{i=1}^{\infty} c_{i, \rho(i)} \end{aligned}$$

where $c_{i,j} = 2(a_{i,j} + b_{i,j}) \forall i, j \in \mathbb{N}$ is a (D) -sequence, then assertion follows. \square

Using Theorem 3.6 and Definition 3.5 above, we can see immediately that the following holds.

Corrolary 3.7 : Given an elementary set $A \subset \mathfrak{R}^n$ and α volume on A . A function $f : A \rightarrow L$ is said to be Henstock-Kurzweil integrable on A with respect to α , denoted by $f \in HK(A, L, \alpha)$, if $f \in HK(A_i, L, \alpha)$ for every i , where $A = \bigcup_{i=1}^p A_i$ and $\{A_1, A_2, \dots, A_p\}$ is any division on A . The Henstock-Kurzweil integral of function f on A is

$$(HK) \int_A f d\alpha = \sum_{i=1}^p \int_{A_i} f d\alpha.$$

We now state version of the Cauchy criterion.

Theorem 3.8 : A function $f : A \rightarrow L$ is Henstock-Kurzweil integrable if and only if there exists a (D) -sequence $(a_{i,j})_{i,j}$ in L such that, for every $\rho \in \square^\square$ we can find a function $\delta : A \rightarrow \mathfrak{R}^+$ and for every δ -fine Perron partition $P_1 = \{(A, \bar{x})\}$ and $P_2 = \{(I, \bar{x})\}$ on A , we have

$$\left| P_1 \sum f(\bar{x})\alpha(I) - P_2 \sum f(\bar{x})\alpha(I) \right| \leq \bigvee_{i=1}^{\infty} a_{i,\rho(i)}$$

Proof : The proof is similar to the one of Theorem 3.1.8, p. 57 of **Muslim** (2003).

We now prove a result about Hentock-Kurzweil integrability on subcells.

Theorem 3.9 : Let α be a volume on a cell $A \subset \mathfrak{R}^n$. If $f \in HK(A, L, \alpha)$, then $f \in HK(B, L, \alpha)$, for every cell $B \subset A$.

Proof : By virtue of Theorem 3.8, there exists a (D) -sequence $(a_{i,j})_{i,j}$ in L such that, for every $\rho \in \square^\square$ we can find a function $\delta : A \rightarrow \mathfrak{R}^+$ and for every δ -fine Perron partition $P_1 = \{(I, x)\}$ and $P_2 = \{(I, x)\}$ on A , we have

$$\left| P_1 \sum f(\bar{x})\alpha(I) - P_2 \sum f(\bar{x})\alpha(I) \right| \leq \bigvee_{i=1}^{\infty} a_{i,\rho(i)}$$

Since cell $B \subset A$, then there exists a collection of finite non-overlapping cells Γ such that $\overline{A \setminus B} = \bigcup_{C \in \Gamma} C$. By virtue of Cousin Lemma, there exists a δ -fine Perron partition P_C on C , for every $C \in \Gamma$. Let δ -fine Perron partition P_B' and P_B'' on B . Put $P_0 = P_B' \cup \left(\bigcup_{C \in \Gamma} P_C \right)$ and $P_0' = P_B'' \cup \left(\bigcup_{C \in \Gamma} C \right)$. Then P_0 and P_0' are δ -fine Perron partition on A . Moreover, we get

$$\begin{aligned} & \left| P_B' \sum f(\bar{x})\alpha(I) - P_B'' \sum f(\bar{x})\alpha(I) \right| \\ &= \left| P_B' \sum f(\bar{x})\alpha(I) + \sum_{C \in \Gamma} P_C \sum f(\bar{x})\alpha(I) - \sum_{C \in \Gamma} P_C \sum f(\bar{x})\alpha(I) - P_B'' \sum f(\bar{x})\alpha(I) \right| \\ &= \left| P_B' \sum f(\bar{x})\alpha(I) + \sum_{C \in \Gamma} P_C \sum f(\bar{x})\alpha(I) - \left\{ P_B'' \sum f(\bar{x})\alpha(I) + \sum_{C \in \Gamma} P_C \sum f(\bar{x})\alpha(I) \right\} \right| \\ &= \left| P_0 \sum f(\bar{x})\alpha(I) - P_0' \sum f(\bar{x})\alpha(I) \right| \\ &\leq \bigvee_{i=1}^{\infty} a_{i,\rho(i)} \end{aligned}$$

Again, from Theorem 3.8 and the last term, it follows that $f \in HK(B, L, \alpha) \sqcup$

By virtue of Theorem 3.9, we define primitif function of Henstock-Kurzweil integrable function f on a cell $A \subset \mathbb{R}^n$ with respect to a volume α as follows.

Definition 3.10 : If $f \in HK(A, L, \alpha)$ and $\mathfrak{I}(A)$ is a collection of all subcells in A , then a function $F: \mathfrak{I}(A) \rightarrow L$ satisfying

$$F(I) = (HK) \int_I f d\alpha \quad \text{and} \quad F(\emptyset) = \bar{0}$$

for every cell $I \in \mathfrak{I}(A)$ is called α -Primitif of Henstock-Kurzweil integrable function f on $\mathfrak{I}(A)$.

REFERENCES

- Boccuto, A., 1998 , *Differential and integral calculus in Riesz Spaces*, Tatra Mountains Math. Publ.,14,133-323.
- Boccuto, A and Riecan, B., 2004 , *On The Henstock-Kurzweil Integral for Riesz-Space-Valued Functions Defined on Unbounded Intervals*, *Check. Math. Journal*, 54,3, 591-607.
- Indrati, C.H., 2002 , *Integral henstock-Kurzweil pada Ruang Euclide R^n* , Disertasi UGM, Yogyakarta.
- Muslim, A., 2003 , *Integral Henstock-Kurzweil Fungsi-Fungsi dari Ruang Euclide R^n ke Ruang Banach ℓ_∞* , Tesis S2 UGM, Yogyakarta.
- Muslim, A dan Soeparna, D., 2002 , *Integral Henstock Fungsi Bernilai Barisan*, *Jurnal Matematika dan Pembelajaran*, UM, Malang.
- Pfeffer, W.F., 1993 , *The Riemann Approach to Integration*, Cambridge University Press.
- Riecan, B., 1989 , *On the Kurzweil Integral for Functions with Values in Ordered Spaces I*, *Acta Math. Univ. Comenian.* 56-57,75-83.
- Riecan, B., 1992 , *On Operator Valued Measures in Lattice ordered Groups*, *Atti. Sem. Mat. Fis. Univ. Modena*, 40, 151-154.
- Riecan, B and Neubrunn, T., 1997 , *Integral, Measure and Ordering*, Kluwer Academic Publishers, Bratislava.
- Riecan, B and Vrabelova, M., 1996 , *On The Kurzweil integral for Functions with Values in Ordered Spaces III*, *Tatra Mountains Math. Publ.*,8, 93-100.
- Zachriwan., 2004 , *Integral Henstock Fungsi bernilai di dalam Ruang Barisan ℓ_p ($1 \leq p < \infty$)*, Tesis S2 UGM.
- Zaanen, A.A., 1997 , *Introduction to Operator Theory in Riesz Spaces*, Springer Verlag.